KabaFusion
Patient-Focused Infusion Therapy

5860 W. Las Positas Blvd | Suite 19 | Pleasanton, CA 94588
OFFICE: 925.605.8981 | FAX: 833.543.0292

Immunoglobulin Prescription Form

Please fax completed order form to 833.543.0292

Demographic Information:

Patient Name _____ **Date of Birth** _____

Home Address _____

City, State, Zip _____

Home Phone _____ **Mobile or Work Phone** _____

Primary Insurance Name _____

Primary Insurance ID _____ **Primary Insurance Group** _____

Insured Name _____ **Insured Date of Birth** _____

Secondary Insurance Name _____ **Insurance ID** _____ **Insurance Group** _____

Secondary Insurance ID _____ **Secondary Insurance Group** _____

Ordering Physician's Name _____

Address _____

City, State, Zip _____

Phone _____ **Fax** _____

NPI _____

Please fax the following information:

□ History and Physical □ Pertinent Lab Work

□ Front & Back copy(s) of patient's insurance card(s)

I authorize KabaFusion and its representatives to act as an agent and initiate and execute any insurance prior authorization process for this prescription, and any future fills of the same prescription for the patient listed above. I understand that I can revoke this designation at any time by providing written notice to KabaFusion.

Physician Signature:_____

Date: _____

Prescription:

☐ **Intravenous Immunoglobulin** ☐ **Subcutaneous Immunoglobulin**

☐ 0.4 gm/kg ☐ 1gm/kg ☐ 2gm/kg ☐ _____ grams

Infuse: ☐ IV daily x _____ day(s); repeat every _____ week(s) x _____ cycles

☐ Other: _____

Infuse _____ grams OR _____ mls using _____ sites _____ time(s) per week

Hydration order: _____ mls NS iv to be infused prior/post IVIG. for _____ months.

☐ Pre-medications: Acetaminophen 650mg PO 30 mins prior to infusion
Diphenhydramine 25mg PO 30 mins prior to infusion

☐ Other Pre-medications: _____

Clinical Information:

Patient Weight: _____ Height: _____ Allergies: _____

☐ IV access [for IVIg patients only]: _____ ☐ Nurse to place PIV prior to therapy

Diagnosis	ICD-10	Diagnosis	ICD-10
Neuromuscular:		**Immune Deficiency:**	
☐ Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)	G61.81	☐ CVID w/ Predominant Immunoregulatory T-Cell Disorders	D83.1
☐ Dermatopolymyositis	M33.90	☐ Combined Immunodeficiency, Unspecified	D81.9
☐ Guillain-Barre Syndrome (GBS)	G61.0	☐ Common Variable Immunodeficiency, Unspecified	D83.9
☐ Multifocal Motor Neuropathy	G61.82	☐ Hereditary Hypogammaglobulinemia	D80.0
☐ Myasthenia Gravis (MG)	G70.0	☐ Immunodeficiency with Increased IgM	D80.5
☐ Myasthenia Gravis with (Acute) Exacerbation	G70.01	☐ Nonfamilial Hypogammaglobulinemia	D80.1
☐ Polymyositis	M33.20	☐ Other combined Immunodeficiencies	D81.89
☐ Relapsing Remitting Multiple Sclerosis (RRMS)	G35	☐ Other Common Variable Immunodeficiencies	D83.9
☐ Stiff Person Syndrome	G25.82	☐ Pemphigoid	L12.0
Other:		☐ Pemphigus	L10.9
☐ Autoimmune Encephalopathy	G04.81	☐ SCID with Low or Normal B-Cell Numbers	D81.2
☐ Idiopathic Thrombocytopenic Purpura	D69.3	☐ SCID with Low T- and B- Cell Numbers	D81.1
☐ Inflammatory Neuropathies	G61.89	☐ Selective deficiency of IgG Subclasses	D80.3
		☐ Specific Antibody Deficiency	D80.6
		☐ Systemic lupus erythematosus (SLE)	M32.9

Please Draw:

□ CBC/diff □ CMP □ IgG w/subclasses 1-4 □ Quant. Ig

☐ _____ ☐ _____ Frequency: _____

PER Anaphylaxis Protocol:

□ Adult – EpiPen 0.3 auto-injector dual pack

□ Pediatric – EpiPen 0.15 auto-injector dual pack

* Administer intramuscularly in the event of ADR*

[May repeat x 1. **Order is valid for 1 year**]. **Use generic if applicable**

Notes:

If applicable, flush intravenous access device per KabaFusion protocol:

Access	NS	Heparin
Peripheral	1-3ml before/after use	10u/ml 1-2mls after last NS flush
Midline, central (non-port), PICC	NS 5-10 mls before/after use; 10mls after blood draw	10 u/ml 3-5mls after last NS flush; 5mls after blood draw
Implanted Port	5-10mls before/after use; 20mls after blood draw	100 u/ml 5mls after last NS flush; 5mls after blood draw
Tunneled	5-10mls before/after use; 20mls after blood draw	10 u/ml 3- mls after last NS flush. 5mls after blood draw
Groshong PICC, Midline	5-10mls before/after use; 10mls after blood draw	NO Heparin needed

CONFIDENTIALITY NOTICE: The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.