

## Immune Deficiency Immunoglobulin Therapy Patient Referral and Prescription Sheet Return Signed RX via Fax to: 877.309.2209

To:		From:				Phone:					
Intake phone: <b>877.309.2207</b>		Fax:		Numb			er of Pages (Including Cover):				
Date:	DOB:			Alle	Allergies:						
Patient Name:			1		Height:		Weight:				
Rx: Intravenous Route  IVIG grams daily for day(s) OR IVIG grams/kilogram daily given over non-consecutive/ consecutive day(s)  Repeat course every week(s) for a total of course(s) Dose will be rounded to nearest vial size.											
Rx: Subcutaneous Route  IG grams each month given as doses OR IG grams times per month. Administer SQIG using sites at a time. Repeat week(s). Ok to round dose to nearest vial size. Refill x 1yr.											
Diagnosis:			ICD-9	ICD-9 ICD-1		Diagnosis: ICD-9 ICD-10					
Common Variable Immunodeficiency with						Selective deficiency of Immunoglobulin M [IgM]			279.02	D80.4	
Predominant Immunoregulatory T-Cell Disorders		ers	279.10	D83.1		Selective deficiency of Immunoglobulin					
☐ Wiskott-Aldrich Syndrome			279.12	D82.0		G [IgG] Subclasses			279.03	D80.3	
Combined Immunodeficiency, Unspecified				D81.9	)	☐ Hereditary Hypogammaglobulinemia			279.04	D80.0	
Severe Combined Immunodeficiency [SCID]			279.2		L	☐ Immunod	eficiency with In	iciency with Increased IgM		D80.5	
with Low T- and B- Cell Numbers				D81.1		Other Common Variable Immunodeficiencies				D83.8	
Severe combined Immunodeficiency [SCID]with Low or Normal B-Cell Numbers				D81.2	2	Common Variable Immunodeficiency, Unspecified			279.06	D83.9	
Selective deficiency of Immunoglobulin A IgA]			279.01	01 D80.2		Other:				503.5	
IV Access Device: Peripheral Central Hydration: Infuse 500 ml of NS. Start at 100ml/hr 30 minutes before the IVIG starts; then run concurrently with IVIG infusion.  Premedication Orders: Refill x 1Year Per KabaFusion recommendation: -ACETAMINOPHEN 650 MG (325mg X 2) orally  Access NS Heparin 100 u/ml											
PRE-IVIG -DIPHENHYDRAMINE 25 MG orally PRE-IVIG			-			Peripheral	1 - 3 ml before/after use		1 - 3 ml after last NS		
□None						Midline, Central (Non-	3 - 5 ml before/after use		3 - 5 ml		
Other premed orders:						Port), PICC	5 - 10 ml after blood draw		after last NS		
Other premed orders:				_	lı	mplanted Port		5 - 10 ml before/after use 0 - 20 ml after blood draw		5 ml after last NS	
□Other premed orders: □Epi-Pen 0.3mg 2-Pak Auto-Injector				-	Groshong Midlir			ore/after use er blood draw			
If needed, a Basic Metabolic Panel (Chem 7) to be drawn prior to first dose. If necessary Patient may have first dose administered in the home by a skilled nurse, unless instructed otherwise. Pharmacy to provide all equipment and medical supplies necessary to administer therapy as appropriate.  Prescriber Signature:  Date  Print Prescriber Name:  NPI#											
Please fax the following information:  ☑ Immunoglobulin order – include dose, route of administration, frequency, duration, and any pre-medications <b>OR</b> use prescription order section above  ☑ Patient demographics – include insurance information. <b>We will obtain authorization</b> unless the insurance dictates otherwise  ☑ H & P <b>OR</b> progress note(s) describing diagnosis and clinical status ☑ Labs – BUN/Creatinine (preferred within last 90 days), Immunoglobulin Panel											
CONFIDENTIALITY NOTICE  The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.											

KabaFusion Infusion Pharmacy | 3500 NW Boca Raton Blvd | Suite 704 | Boca Raton, FL 33431 Phone: 877.309.2207 | Fax: 877.309.2209 | www.kabafusion.com