

Cutaquig SCIG Therapy Patient Referral and Prescription

| To: | | From: | | | | Phone | : | |
|---|------|-------|-------|---------------------------------------|------------------------------------|-------|---------|--------------|
| Intake phone: 757.424.4822 | | Fax: | | | Number of Pages (Including Cover): | | | ling Cover): |
| Date: | DOB: | I | Aller | gies: | | | | |
| Patient Name: | | | Heigh | t: | | | Weight: | |
| Begin Cutaquig SCIG per KabaFusion protocol formonths Begin Cutaquiggrams SCIG everyformonths KabaFusion to provide infusion pump needle administration sets (A4221) KabaFusion to provide infusion supplies for infusion pump (K0552) KabaFusion to provide mechanical ambulatory infusion pump (E0779) Infusion RN home visit prior to therapy start to assess patient and home environment, educate patient about SCIG therapy. Infusion RN visits x 3 and PRN to educate patient to self administer Cutaquig infusion KabaFusion to provide all professional services related to infusion | | | | | | | | |
| Diagnosis: | | | | | | | | ICD-10 |
| Common Variable Immunodeficiency with Predominant Immunoregulatory T-Cell Disorders | | | | | | | | D83.1 |
| Wiskott-Aldrich Syndrome | | | | | | | | D82.0 |
| Combined Immunodeficiency, Unspecified | | | | | | | | D81.9 |
| Severe Combined Immunodeficiency [SCID] with Low T- and B- Cell Numbers | | | | | | | D81.1 | |
| Severe combined Immunodeficiency [SCID] with Low or Normal B-Cell Numbers | | | | | | | | D81.2 |
| Selective deficiency of Immunoglobulin A IgA] | | | | | | | | D80.2 |
| Selective deficiency of Immunoglobulin M [IgM] | | | | | | | | D80.4 |
| Selective deficiency of Immunoglobulin G [IgG] Subclasses | | | | | | | D80.3 | |
| Hereditary Hypogammaglobulinemia | | | | | | | D80.0 | |
| Immunodeficiency with Increased IgM | | | | | | | D80.5 | |
| Other Common Variable Immunodeficiencies | | | | | | | | D83.8 |
| Common Variable Immunodeficiency, Unspecified | | | | | | | | D83.9 |
| Other: | | | | | | | | |
| Premedication Orders: | | | | DIPHENHYDRAMINE 25 MG orally PRE-SCIG | | | | |
| Refill x 1Year Other: Per KabaFusion recommendation: Epinephrine 0.3mg 2-Pak Auto-Injector ACETAMINOPHEN 650 MG (325mg X 2) orally PRE-SCIG None | | | | | | | DL | |
| Prescriber Signature:Date | | | | | | | | |
| Print Prescriber Name:NPI# | | | | | | | | |
| Please fax the following information: Immunoglobulin order – include dose, route of administration, frequency, duration, and any pre-medications OR use prescription order section above Patient demographics – include insurance information. We will obtain authorization unless the insurance dictates otherwise H & P OR progress note(s) describing diagnosis and clinical status Labs – BUN/Creatinine (preferred within last 90 days), ImmunoglobulinPanel | | | | | | | | |
| document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you. KabaFusion Infusion Pharmacy 816 Greenbrier Cir. Suite E Chesapeake, VA 23320 | | | | | | | | |
| Phone: 757.424.4822 Fax: 757.424.5871 <u>www.kabafusion.com</u> | | | | | | | | |