

## Remicade (infliximab) Patient Referral and Prescription Sheet Return Signed Rx via Fax to: 704.583.2130

Date: To:		From:		
Phone: <b>866.583.2140</b>		Fax:	Numbe	er of Pages:
		Patient Information		
Patient Name: DOB:				Height:
Allergies:				Weight:
Medication Order				
☐ Remicade® (infliximab) ☐ Inflectra® (infliximab-dyyb) ☐ Renflexis® (infliximab-abda) ☐ Avsola™ (infliximab-axxq)				
	innectra® (innixima	ab-dyyb) <b>i kennexis</b>		Avsola (IIIIIXIIIIab-axxq)
□kg	= IV at		eeks for cours	es
Dose will be rounded to the n Reconstitute each vial w Administer for at least 2 hours	vith 10 mL Sterile Wate	er then dilute with 250	) mL Normal Saline.	Refills
Diagnosis	ICD-10		Diagnosis	ICD-10
Ankylosing spondylitis	M45	☐ Rheumatoid arthritis		M06.9
Crohn's Disease	K50.90	☐ Ulcerative colitis K5		K51.90
Plaque psoriasis	L40.0	Other:		
☐ Psoriatic arthritis	L40.52			
IV Access Device: ☐ Perip	heral 🗖 Central	Lab order: 🗖 CBC with	h diff	
Pre-medications: If applicable, flush IV access device per KabaFusion protoc				sion protocol:
☐ Diphenhydramine: ☐ PO ☐ IV		Access	NS	Heparin
☐ Acetaminophen: ☐ PO ☐ IV		Peripheral	1-3 mL before/after use	10 U/mL 1-2 mL after last NS flush
☐ Methylprednisolone: ☐ IV ☐ Other pre-meds:		Midline, Central (non-port), PICC	5-10 ml before/after use 5-20 ml after blood draw	10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw
☐ <b>Hydration:</b> Infuse 500 mL of Normal Saline with Infliximab infusion		Implanted Port	5-10 mL before/after use.	100 U/mL 5 mL after last NS flush
☐ Anaphylaxis Protocol: Epinephrine Auto-Injector dual pack:			20mL after blood draw	5 mL after the blood draw
Adult: 0.3 mg <u>Children</u> : 0.15 mg		Tunneled	5-10 mL before/after use 20mL after blood draw	10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw
Administer epinephrine IM in the event of anaphylaxis. May repeat x 1 as needed, Call 911. <b>Refill x 1yr</b>		Groshong PICC, Midline	5-10 mL before/after use	None
Lauthorize KahaFusion and its ren	resentatives to act as an agent	and initiate and execute any	10 mL after blood draw	on process for this prescription and any
I			•	providing written notice to KabaFusion.
			, ,	
Prescriber Signature:			Date:	
Print Prescriber Name:NPI#:				
	DOCUMENTA	ATION – PLEASE FAX TO	K ABAFUS ION	
□ Rx Order – include dose, route of administration, frequency, duration, & pre-medications OR use the Rx order form above				
Patient Demographics – include <u>insurance</u> information. <u>We will obtain authorization</u> unless the insurance dictates otherwise.				
☐ H&POR progress note(s) de				
□ TB and Hepatitis B Virus (HBV) screening results (required prior to Remicade initiation) – HBsAg and anti-HBc. □ Most recent lab results for: BUN/Creatinine (preferred within last 90 days). CMP Panel and CBC with Differential				

## CONFIDENTIALITY NOTICE

The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.