



Rituxan (Rituximab) Patient Referral and Prescription Sheet

Return Signed Rx via Fax to: 888.837.2716

Date:	To:	From:																			
Phone:	Fax:	Number of Pages:																			
Patient Information																					
Patient Name:	DOB:	Height:																			
Allergies:			Weight:																		
Medication Order																					
Rituximab: <input type="checkbox"/> Rituximab: 375mg x _____ BSA (m ²) = _____ IV every _____ for _____ courses. <input type="checkbox"/> Rituximab: _____ -First Rituximab IV dose is to be administered in a controlled environment setting then subsequent doses in the home setting -Dilute Rituxan to a final concentration of 1 mg/mL - 4 mg/mL with either 0.9% Sodium Chloride, or 5% Dextrose			Refills																		
Diagnosis	ICD-10	Diagnosis	ICD-10																		
<input type="checkbox"/> Rheumatoid Arthritis	M06. 9	<input type="checkbox"/> Wegener's Granulomatosis	M31.3																		
<input type="checkbox"/> Chronic Lymphocytic Leukemia (CLL)	C91.10	<input type="checkbox"/> Pemphigus Vulgaris (PV)	L10.0																		
<input type="checkbox"/> Non-Hodgkin's Lymphoma (NHL)	C85. 9	Other:																			
<input type="checkbox"/> Neuromyelitis Optica [Devic]	G36.0																				
IV Access Device: <input type="checkbox"/> Peripheral <input type="checkbox"/> Central		Lab order: <input type="checkbox"/> CBC with diff <input type="checkbox"/> CMP <input type="checkbox"/> _____																			
Pre-medications: <input type="checkbox"/> Diphenhydramine: _____ <input type="checkbox"/> PO <input type="checkbox"/> IV <input type="checkbox"/> Acetaminophen: _____ <input type="checkbox"/> PO <input type="checkbox"/> IV <input type="checkbox"/> Methylprednisolone: _____ <input type="checkbox"/> IV <input type="checkbox"/> Other pre-meds: _____ <input type="checkbox"/> Hydration: Infuse 500 mL of Normal Saline with Rituximab infusion <input type="checkbox"/> Anaphylaxis Protocol: Epinephrine Auto-Injector dual pack: <u>Adult:</u> 0.3 mg <u>Children:</u> 0.15 mg Administer epinephrine IM in the event of anaphylaxis. May repeat x 1 as needed, Call 911. Refill x 1yr		If applicable, flush IV access device per KabaFusion protocol: <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 25%;">Access</th> <th style="width: 25%;">NS</th> <th style="width: 50%;">Heparin</th> </tr> </thead> <tbody> <tr> <td>Peripheral</td> <td>1-3 mL before/after use</td> <td>10 U/mL 1-2 mL after last NS flush</td> </tr> <tr> <td>Midline, Central (non-port), PICC</td> <td>5-10 ml before/after use 5-20 ml after blood draw</td> <td>10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw</td> </tr> <tr> <td>Implanted Port</td> <td>5-10 mL before/after use. 20mL after blood draw</td> <td>100 U/mL 5 mL after last NS flush 5 mL after the blood draw</td> </tr> <tr> <td>Tunneled</td> <td>5-10 mL before/after use 20mL after blood draw</td> <td>10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw</td> </tr> <tr> <td>Groshong PICC, Midline</td> <td>5-10 mL before/after use 10 mL after blood draw</td> <td>None</td> </tr> </tbody> </table>		Access	NS	Heparin	Peripheral	1-3 mL before/after use	10 U/mL 1-2 mL after last NS flush	Midline, Central (non-port), PICC	5-10 ml before/after use 5-20 ml after blood draw	10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw	Implanted Port	5-10 mL before/after use. 20mL after blood draw	100 U/mL 5 mL after last NS flush 5 mL after the blood draw	Tunneled	5-10 mL before/after use 20mL after blood draw	10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw	Groshong PICC, Midline	5-10 mL before/after use 10 mL after blood draw	None
Access	NS	Heparin																			
Peripheral	1-3 mL before/after use	10 U/mL 1-2 mL after last NS flush																			
Midline, Central (non-port), PICC	5-10 ml before/after use 5-20 ml after blood draw	10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw																			
Implanted Port	5-10 mL before/after use. 20mL after blood draw	100 U/mL 5 mL after last NS flush 5 mL after the blood draw																			
Tunneled	5-10 mL before/after use 20mL after blood draw	10 U/mL 3-5 mL after last NS flush 5 mL after the blood draw																			
Groshong PICC, Midline	5-10 mL before/after use 10 mL after blood draw	None																			
I authorize KabaFusion and its representatives to act as an agent and initiate and execute any insurance prior authorization process for this prescription and any future fills of the same prescription for the patient listed above. I understand that I can revoke this designation at any time by providing written notice to KabaFusion.																					
Prescriber Signature: _____		Date: _____																			
Print Prescriber Name: _____		NPI#: _____																			
DOCUMENTATION – PLEASE FAX TO KABAFUSION																					
<input type="checkbox"/> Rx Order – include dose, route of administration, frequency, duration, & pre-medications OR use the Rx order form above <input type="checkbox"/> Patient Demographics – include insurance information. <u>We will obtain authorization</u> unless the insurance dictates otherwise. <input type="checkbox"/> H & P OR progress note(s) describing diagnosis, clinical status, and clinical symptoms. <input type="checkbox"/> TB and Hepatitis B Virus (HBV) screening results (required prior to Rituximab initiation) – HBsAg and anti-HBc. <input type="checkbox"/> Most recent lab results for: BUN/Creatinine (preferred within last 90 days), CMP Panel and CBC with Differential																					

CONFIDENTIALITY NOTICE

The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.