

SIMPONI ARIA (GOLIMUMAB) PATIENT REFERRAL AND PRESCRIPTION SHEET

	Pati	ient Information				
Patient Name:			Date:			
DOB:			Height:	🗆 inches 🗆	🗅 inches 🕒 cm	
Allergies:			Weight:	🗆 lbs 🖵 kg	🗆 lbs 🖵 kg	
Primary Diagnosis						
Diagnosis ICD-10 Diagnosis			ICD-10		ICD-10	
Rheumatoid arthritis	M45	Rheumatoid arthritis with rheumatoid factor, unspecified		M05.9		
Rheumatoid arthritis, unspecified	M06.9	Rheumatoid arthritis w/o rheumatoid factor, unspecified		M06.00		
Juvenile rheumatoid polyarthritis (seronegative)	M08.3	Other juvenile arthrit	Other juvenile arthritis, unspecified site		M08.80	
Psoriatic Arthritis	L40.52	Other:			- -	
Ankylosing Spondylitis	M45					
Medication Order						
Simponi Aria (Golimumab): 2mg/kg IV at weeks 0,4 and then every 8 weeks x 1 year (initial dosing) 2mg/kg IV every 8 weeks x 1 year Other			from date signature indicated	 Refills x one year from date of signature unless indicated below Refills 		
Ancillary Orders						
Pre-medications:	IV Eluch Ordore	 IV Flush Orders: Peripheral: NS 1-3 mL before/after use Implanted VAD: NS 5 to 10 mL before/after use and 10 mL post-lab draw. Heparin (100 unit/mL) 3 to 5 mL final flush CVAD: NS 5 to 10 mL before/after use and 10 mL post-lab draw Heparin (10 units/mL) 3 to 5 mL final flush Refill x one year 				
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds: Refill x one year 		 Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 un) 	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) 10 mL before/after use nits/mL) 3 to 5 mL final f	′after use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr	aw	
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds:	•	 Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 ur Refill x one yee 0.3mg Children: 0.15 mg 	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) = 10 mL before/after use nits/mL) 3 to 5 mL final f ear	′after use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr	aw	
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds: Refill x one year Anaphylaxis Protocol: Epinephrine Auto-Injector dual 	•	 Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 ur Refill x one yee 0.3mg Children: 0.15 mg 	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) = 10 mL before/after use nits/mL) 3 to 5 mL final f ear	Yafter use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr flush	aw	
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds: Refill x one year Anaphylaxis Protocol: Epinephrine Auto-Injector dual Administer epinephrine IM in the event of anaphylaxis. 	May repeat >	 Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 un Refill x one yet 0.3mg Children: 0.15 mg x 1 as needed, Call 911. F sults Patient Demogobtain authori H&P OR progr Medication Lis 	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) : 10 mL before/after use nits/mL) 3 to 5 mL final f ar Refill x 1yr Other Required Do graphics – include insura ization unless the insura	Yafter use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr flush cumentation ance information. We nce dictates otherwise resent DMARDS and	will	
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds: Refill x one year Anaphylaxis Protocol: Epinephrine Auto-Injector dual Administer epinephrine IM in the event of anaphylaxis. Therapy Specific Documentation Please include the following lab results required for infu are available, the following labs will be drawn prior to fi Hepatitis B Surface Antigen Hepatitis B Core Antibody Total (not Core IgM) QuantiFERON TB 	May repeat >	 Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 un Refill x one yet 0.3mg Children: 0.15 mg x 1 as needed, Call 911. F sults Patient Demogobtain authori H&P OR progr Medication Lis 	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) : 10 mL before/after use nits/mL) 3 to 5 mL final f ar Gar Other Required Do graphics – include insura ization unless the insura ess note(s) st - please list past and p	Yafter use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr flush cumentation ance information. We nce dictates otherwise resent DMARDS and	will	
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds: Refill x one year Anaphylaxis Protocol: Epinephrine Auto-Injector dual Administer epinephrine IM in the event of anaphylaxis. Therapy Specific Documentation Please include the following lab results required for infu are available, the following labs will be drawn prior to fi Hepatitis B Surface Antigen Hepatitis B Core Antibody Total (not Core IgM) QuantiFERON TB 	May repeat >	 Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 ur Refill x one yee 0.3mg Children: 0.15 mg x 1 as needed, Call 911. F sults Patient Demog obtain authori H&P OR progr Medication Lis biologics below 	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) : 10 mL before/after use nits/mL) 3 to 5 mL final f ar Gar Other Required Do graphics – include insura ization unless the insura ess note(s) st - please list past and p	Yafter use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr flush cumentation ance information. We nce dictates otherwise resent DMARDS and	will	
 Diphenhydramine: 25mg PO 30 min pre-infusion Acetaminophen: 650mg PO 30 min pre-infusion Famotidine: 20 mg PO x1 dose Other pre-meds: Refill x one year Anaphylaxis Protocol: Epinephrine Auto-Injector dual Administer epinephrine IM in the event of anaphylaxis. Therapy Specific Documentation Please include the following lab results required for infu are available, the following labs will be drawn prior to fit Hepatitis B Surface Antigen Hepatitis B Core Antibody Total (not Core IgM) QuantiFERON TB Other: 	May repeat >	Peripheral: NS Implanted VAI post-lab draw. CVAD: NS 5 to Heparin (10 ur Refill x one ye 0.3mg Children: 0.15 mg x 1 as needed, Call 911. F sults Patient Demog obtain authori H&P OR progr Medication Lis biologics below //der Information Pro	D: NS 5 to 10 mL before/ Heparin (100 unit/mL) = 10 mL before/after use nits/mL) 3 to 5 mL final f ear Other Required Do graphics – include insura ization unless the insura ess note(s) st - please list past and p w with dates of discontin	Yafter use and 10 mL 3 to 5 mL final flush and 10 mL post-lab dr flush cumentation ance information. We nce dictates otherwise resent DMARDS and	will	

I authorize KabaFusion and its representatives to act as an agent and initiate and execute any insurance prior authorization process for this prescription and any future refills of the same prescription for the patient listed above. I understand that I can revoke this designation at any time by providing written notice to KabaFusion.

Prescriber Signature: ____

Date: _____

CONFIDENTIALITY NOTICE: The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.