



Prescriber Signature: _

ULTOMIRIS (RAVULIZUMAB-CWVZ) PATIENT REFERRAL AND PRESCRIPTION SHEET					
Patient Information					
Patient Name:			Date:		
DOB:			Height:	Height:	
Allergies:			Weight:		□ lbs □ kg
Primary Diagnosis					
Diagnosis ICD-10 Diagnosis ICD-10					
☐ Paroxysmal Nocturnal Hemoglobinuria	D59.5	☐ Generalized Myast	henia Gravis		G70.00
☐ Atypical Hemolytic Uremic Syndrome	D59.39	☐ Neuromyelitis Opti	ica Spectrum Disorder		G36.0
☐ Other:					
Medication Order					
Loading Dose ☐ Infuse 2400mg IV x 1 dose (pt weight 40-59kg) ☐ Infuse 2700mg IV x 1 dose (pt weight 60-99kg) ☐ Infuse 3000mg IV x 1 dose (pt weight ≥100kg) ☐ Other Maintenance Dose ☐ Infuse 3000 mg IV every 8 weeks after loading dose (pt weight 40-59kg) ☐ Infuse 3300 mg IV every 8 weeks after loading dose (pt weight 60-99kg) ☐ Infuse 3600mg IV every 8 weeks after loading dose (pt weight ≥100kg) ☐ Other				□ Refills x one year from date of signature unless indicated below □ Refills	
Nursing Orders Skilled nursing visit for clinical assessment, administration of medication. Initiate plan of treatment for ongoing nursing services. Insert or access and maintain access device as indicated in Ancillary Orders. Remove peripheral IV or access from implanted VAD when infusion is completed.					
Ancillary Orders					
		Ancillary Orders			
Pre-medications: ☐ Diphenhydramine: 25mg PO 30 min pre-infus ☐ Acetaminophen: 650mg PO 30 min pre-infusion ☐ Famotidine: 20 mg PO x1 dose ☐ Other pre-meds: ☐ Refill x one year	ion	IV Flush Ordo Peripheral Implanted lab draw. I	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use L0 units/mL) 3 to 5 mL final	/after use 5 mL final and 10 m	flush
☐ Diphenhydramine: 25mg PO 30 min pre-infus ☐ Acetaminophen: 650mg PO 30 min pre-infusi ☐ Famotidine: 20 mg PO x1 dose ☐ Other pre-meds:	ion on ctor dual pack: Adul	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use L0 units/mL) 3 to 5 mL final e year	/after use 5 mL final and 10 m	flush
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusi □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje	ion on ctor dual pack: Adul ophylaxis. May repea	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use L0 units/mL) 3 to 5 mL final e year	/after use 5 mL final and 10 m flush	flush IL post-lab draw
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Injee Administer epinephrine IM in the event of analysis.	ion on ctor dual pack: Adul aphylaxis. May repea tation	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1 Refill x on t: 0.3mg Children: 0.19 t x 1 as needed, Call 9 Patient De obtain aut H&P OR p	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Do emographics – include insurationization unless the insurar orgress note(s) n List (include prior/failed E	/after use 5 mL final e and 10 m flush cocumenta ance infor	flush nL post-lab draw tion mation. We will tes otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusi □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen Please include the following lab results □ MuSK Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mer	ion ctor dual pack: Adul phylaxis. May repea	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1 Refill x on t: 0.3mg Children: 0.19 t x 1 as needed, Call 9 Patient De obtain aut H&P OR p Medicatio	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Do emographics – include insurationization unless the insurar orgress note(s) n List (include prior/failed E	/after use 5 mL final e and 10 m flush cocumenta ance infor	flush nL post-lab draw tion mation. We will tes otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusi □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen Please include the following lab results □ MuSK Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mer	ion ctor dual pack: Adul phylaxis. May repea	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1 Refill x on t: 0.3mg Children: 0.15 t x 1 as needed, Call 9 Patient De obtain aut H&P OR p Medicatio steroid us	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Do emographics – include insurationization unless the insurar orgress note(s) n List (include prior/failed E	/after use 5 mL final e and 10 m flush cocumenta ance infor	flush nL post-lab draw tion mation. We will tes otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusi □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen Please include the following lab results □ MuSK Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mer	ion ctor dual pack: Adul phylaxis. May repea	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1 Refill x on t: 0.3mg Children: 0.15 t x 1 as needed, Call 9 Patient De obtain aut H&P OR p Medicatio steroid us	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Doministry of the comparable of	/after use 5 mL final e and 10 m flush cocumenta ance infor	flush nL post-lab draw tion mation. We will tes otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infusion □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inject Administer epinephrine IM in the event of anatomister epinephrine IM in	ctor dual pack: Adul aphylaxis. May repea tation	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1 Refill x on t: 0.3mg Children: 0.19 t x 1 as needed, Call 9 Patient De obtain aut H&P OR p Medicatio steroid use	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before 10 units/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Does mographics – include insuration unless the insuration gress note(s) in List (include prior/failed Ee) Provider Phone: Provider Fax:	/after use 5 mL final e and 10 m flush cocumenta ance infor	flush nL post-lab draw tion mation. We will tes otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusi □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen Please include the following lab results □ MuSK Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mer □ Other □ Other □ Provider Name: Provider NPI:	ctor dual pack: Adul aphylaxis. May repea tation	IV Flush Ord Peripheral Implanted lab draw. CVAD: NS Heparin (1 Refill x on t: 0.3mg Children: 0.19 t x 1 as needed, Call 9 Patient De obtain aut H&P OR p Medicatio steroid use	l: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before 10 units/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Does mographics – include insuration unless the insuration gress note(s) in List (include prior/failed Ee) Provider Phone: Provider Fax:	/after use 5 mL final e and 10 m flush cocumenta ance infor	flush nL post-lab draw tion mation. We will tes otherwise.

CONFIDENTIALITY NOTICE: The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.

Date: _