



Prescriber Signature: _

ULTOMIRIS (RAVULIZUMAB-CWVZ) PATIENT REFERRAL AND PRESCRIPTION SHEET

ODIOMINIS (NAVODIZOMAD GWVZ) TATIENT KEI ERRAD AND TRESCRIT HON SHEET					
Patient Information					
Patient Name: Date:					
DOB:			Height:	Height:	
Allergies:			Weight:		lbs 🗆 kg
Primary Diagnosis					
Diagnosis	ICD-10	Diagnosis	osis ICD-10		
☐ Paroxysmal Nocturnal Hemoglobinuria	D59.5	☐ Generalized Myast	ieneralized Myasthenia Gravis		G70.00
☐ Atypical Hemolytic Uremic Syndrome	D59.39	☐ Neuromyelitis Opti	ca Spectrum Disorder		G36.0
☐ Other:					
Medication Order					
Loading Dose ☐ Infuse 2400mg IV x 1 dose (pt weight 40-59kg) ☐ Infuse 2700mg IV x 1 dose (pt weight 60-99kg) ☐ Infuse 3000mg IV x 1 dose (pt weight ≥100kg) ☐ Other Maintenance Dose ☐ Infuse 3000 mg IV every 8 weeks after loading dose (pt weight 40-59kg) ☐ Infuse 3300 mg IV every 8 weeks after loading dose (pt weight 60-99kg) ☐ Infuse 3600mg IV every 8 weeks after loading dose (pt weight ≥100kg) ☐ Other Nursing Orders Skilled nursing visit for clinical assessment, administration of medication. Initiate plan of tr					
maintain access device as indicated in Ancillary Orders. Remove peripheral IV or access from implanted VAD when infusion is completed.					
Ancillary Orders					
Pre-medications: ☐ Diphenhydramine: 25mg PO 30 min pre-infusion ☐ Acetaminophen: 650mg PO 30 min pre-infusion ☐ Famotidine: 20 mg PO x1 dose ☐ Other pre-meds: ☐ Refill x one year					
☐ Diphenhydramine: 25mg PO 30 min pre-infus ☐ Acetaminophen: 650mg PO 30 min pre-infusio ☐ Famotidine: 20 mg PO x1 dose ☐ Other pre-meds:	on	☐ Implanted lab draw. ☐ CVAD: NS	: NS 1-3 mL before/after us VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use L0 units/mL) 3 to 5 mL final	/after use a 5 mL final f and 10 ml	flush
☐ Diphenhydramine: 25mg PO 30 min pre-infus ☐ Acetaminophen: 650mg PO 30 min pre-infusio ☐ Famotidine: 20 mg PO x1 dose ☐ Other pre-meds:	on ctor dual pack: Adul	☐ Peripheral ☐ Implanted lab draw. ☐ CVAD: NS Heparin (1 ☐ Refill x on t: 0.3mg Children: 0.15	I: NS 1-3 mL before/after us VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use L0 units/mL) 3 to 5 mL final e year	/after use a 5 mL final f and 10 ml	flush
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusio □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje	c tor dual pack: Adul phylaxis. May repea	☐ Peripheral ☐ Implanted lab draw. ☐ CVAD: NS Heparin (1 ☐ Refill x on t: 0.3mg Children: 0.15	I: NS 1-3 mL before/after us VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use L0 units/mL) 3 to 5 mL final e year	/after use a 5 mL final f e and 10 ml flush	ilush L post-lab draw
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusio □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana	c tor dual pack: Adul phylaxis. May repea	☐ Peripheral ☐ Implanted lab draw. I ☐ CVAD: NS Heparin (1 ☐ Refill x on It: 0.3mg Children: 0.15 It x 1 as needed, Call 9	I: NS 1-3 mL before/after us VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr	/after use a 5 mL final f e and 10 ml flush	iush L post-lab draw
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusio □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen	ctor dual pack: Adul phylaxis. May repea tation	□ Peripheral □ Implanted lab draw. I □ CVAD: NS Heparin (1 □ Refill x on It: 0.3mg Children: 0.19 It x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p	I: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final 10 e year 11. Refill x 1yr Other Required Decemographics – include insuration unless the insurar orgress note(s) 11. List (include prior/failed E	/after use a 5 mL final f e and 10 ml flush cocumentat rance inforr	ion mation. We will es otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen Please include the following lab results □ MuSk Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mer	ctor dual pack: Adul phylaxis. May repea tation	□ Peripheral □ Implanted lab draw. I □ CVAD: NS Heparin (1 □ Refill x on it: 0.3mg Children: 0.19 it x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p □ Medicatio	I: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final 10 e year 11. Refill x 1yr Other Required Decemographics – include insuration unless the insurar orgress note(s) 11. List (include prior/failed E	/after use a 5 mL final f e and 10 ml flush cocumentat rance inforr	ion mation. We will es otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infus □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Inje Administer epinephrine IM in the event of ana Therapy Specific Documen Please include the following lab results □ MuSk Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mer	ctor dual pack: Adul phylaxis. May repea tation	□ Peripheral □ Implanted lab draw. □ □ CVAD: NS Heparin (1 □ Refill x on it: 0.3mg Children: 0.19 it x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p □ Medicatio steroid us	I: NS 1-3 mL before/after us I VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final 10 e year 11. Refill x 1yr Other Required Decemographics – include insuration unless the insurar orgress note(s) 11. List (include prior/failed E	/after use a 5 mL final f e and 10 ml flush cocumentat rance inforr	ion mation. We will es otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infusion □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Injetadminister epinephrine IM in the event of anatomister epinephrine IM in th	ctor dual pack: Adul phylaxis. May repea tation	□ Peripheral □ Implanted lab draw. □ □ CVAD: NS Heparin (1 □ Refill x on it: 0.3mg Children: 0.19 it x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p □ Medicatio steroid us	i: NS 1-3 mL before/after us VAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Demographics – include insuration unless the insurar orgress note(s) n List (include prior/failed Ee)	/after use a 5 mL final f e and 10 ml flush cocumentat rance inforr	ion mation. We will es otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infusion □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Injete Administer epinephrine IM in the event of anatomic Therapy Specific Document Please include the following lab results □ MuSK Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Menual Country of the provider Name: □ Provider Name:	ctor dual pack: Adul phylaxis. May repea tation	□ Peripheral □ Implanted lab draw. □ CVAD: NS Heparin (1 □ Refill x on it: 0.3mg Children: 0.1! at x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p □ Medicatio steroid us	I: NS 1-3 mL before/after us IVAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Demographics – include insuration unless the insurar orgress note(s) n List (include prior/failed Ee) Provider Phone: Provider Fax:	/after use a 5 mL final f e and 10 ml flush cocumentat rance inforr	ion mation. We will es otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infusion □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Injetadminister epinephrine IM in the event of anatomister epinephrine IM in th	ctor dual pack: Adul phylaxis. May repea tation	□ Peripheral □ Implanted lab draw. □ CVAD: NS Heparin (1 □ Refill x on it: 0.3mg Children: 0.1! at x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p □ Medicatio steroid us	I: NS 1-3 mL before/after us IVAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Demographics – include insuration unless the insurar orgress note(s) n List (include prior/failed Ee) Provider Phone: Provider Fax:	/after use a 5 mL final f e and 10 ml flush cocumentat rance inforr	ion mation. We will es otherwise.
□ Diphenhydramine: 25mg PO 30 min pre-infusion □ Acetaminophen: 650mg PO 30 min pre-infusion □ Famotidine: 20 mg PO x1 dose □ Other pre-meds: □ Refill x one year □ Anaphylaxis Protocol: Epinephrine Auto-Injete Administer epinephrine IM in the event of anatomic Therapy Specific Document Please include the following lab results □ MuSK Test □ ACHR Test □ MG-ADL Score □ Meningococcal Vaccines (MenACWY and Mening Other □ Other □ Other □ Other □ Provider Name: Provider Name: Provider NPI: Is the provider enrolled in Ultomiris REMS Programments of the provider Progr	ctor dual pack: Adul phylaxis. May repea tation B) Pr ram? If not, please	□ Peripheral □ Implanted lab draw. I □ CVAD: NS Heparin (1 □ Refill x on t: 0.3mg Children: 0.1! at x 1 as needed, Call 9 □ Patient De obtain aut □ H&P OR p □ Medicatio steroid us	I: NS 1-3 mL before/after us IVAD: NS 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before Heparin (100 unit/mL) 3 to 5 to 10 mL before/after use 10 units/mL) 3 to 5 mL final e year 5 mg 11. Refill x 1yr Other Required Demographics – include insuration unless the insuration corresponds to 1 include prior/failed Eep Provider Phone: Provider Phone: Provider Fax: rems.com	/after use a 5 mL final fe and 10 ml flush ocumentate ance informance dictate DMARDS, b	ion mation. We will es otherwise. iologics, or

CONFIDENTIALITY NOTICE: The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.

Date: